

AD-A128 420


ON THE TEMPERATURE DEPENDENCE OF THE FORMATION CONSTANT
OF THIOCYANATOPEN. (U) ARMY ARMAMENT RESEARCH AND
DEVELOPMENT COMMAND ABERDEEN PROVI. J R WARD APR 83
ARCSL-TR-82089


1/1

UNCLASSIFIED

F/G 7/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

13

AD

AD A 128420



**CHEMICAL
SYSTEMS
LABORATORY**

US Army Armament Research and Development Command
Aberdeen Proving Ground, Maryland 21010

TECHNICAL REPORT ARCSL-TR-82089

**ON THE TEMPERATURE DEPENDENCE OF THE FORMATION
OF THIOCYANATOPENTAAQUOCHROMIUM (III)
IN ACIDIC SOLUTION**

by

J. Richard Ward

**Chemical Branch
Research Division**

April 1983



[Handwritten signature]
A

Approved for public release; distribution unlimited.

33 05 23 188

DTIC FILE COPY

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|-------------------------------------|---|
| 1. REPORT NUMBER ARCSL-TR-82089 | 2. GOVT ACCESSION NO. AD-A128478 | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) ON THE TEMPERATURE DEPENDENCE OF THE FORMATION CONSTANT OF THIOCYANATOPENTAAQUOCHROMIUM (III) IN ACIDIC SOLUTION | | 5. TYPE OF REPORT & PERIOD COVERED Technical Report |
| 7. AUTHOR(s) J. Richard Ward | | 6. PERFORMING ORG. REPORT NUMBER |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS Commander, Chemical Systems Laboratory ATTN: DRDAR-CLB-CA Aberdeen Proving Ground, Maryland 21010 | | 8. CONTRACT OR GRANT NUMBER(s) |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Commander, Chemical Systems Laboratory ATTN: DRDAR-CLJ-R Aberdeen Proving Ground, Maryland 21010 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1L161101A71A Research In Defense Systems |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 12. REPORT DATE April 1983 |
| | | 13. NUMBER OF PAGES 34 |
| | | 15. SECURITY CLASS. (of this report) UNCLASSIFIED |
| | | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Formation constants for metal ion complexes Blandamer's method Chromium (III) complexes Heat capacity of activation Equilibrium and kinetics of reactions in solution | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The equilibrium constants for the ionization of carboxylic acids in water pass through a maximum as temperature changes. If the equilibrium constant represents a one-step process, then the attendant thermodynamic properties such as enthalpy and entropy are also temperature dependent. Blandamer and co-workers in the United Kingdom and Canada have recently concluded that the ionization of the carboxylic acids is a multistage process and the temperature dependence of the observed equilibrium (Continued). | | |

DD FORM 1473

JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20 ABSTRACT (Cont'd)

constant can be fit by combination of the equilibrium constants of the individual steps, each such equilibrium constant having temperature-independent enthalpy. Blandamer and his co-workers also derived an expression to test the temperature dependence of the enthalpy. <

Temperature dependent enthalpies of reaction are less common in inorganic systems. In 1955, however, Postmus and King noted that the equilibrium constant for the association of thiocyanatopentaaquochromium (III), $\text{Cr}[\text{H}_2\text{O}]_5\text{NCS}^{2+}$, passed through a minimum temperature over the temperature range from 14° to 95° C. Blandamer's equation is used in this report to test the temperature dependence of the heat capacity. Application of Blandamer's equation to Postmus and King's data revealed that the heat capacity was temperature-independent (-66 cal/mole-K) implying that the equilibrium constant for association of $\text{Cr}[\text{H}_2\text{O}]_5\text{NCS}^{2+}$ is a unitary process. This suggests that many other equilibrium and kinetic constants for inorganic ions should also show this temperature behavior, and failure to see it rests on the limited temperature range over which most measurements are made. This also suggests that it is dangerous to extrapolate equilibrium or rate data far outside the range of measurements.

PREFACE

This work was performed under Project 1L161101A71A, Research in Defense Systems, Scientific Area B, during the author's residence at CSL in FY82 as part of cooperative program between BRL and the Research Division at CSL.

The use of trade names in this report does not constitute an official endorsement or approval of the use of such commercial hardware or software. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, Chemical Systems Laboratory, ATTN: DRDAR-CLJ-R, Aberdeen Proving Ground, Maryland 21010. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for United States Government purposes.



| | |
|-----------------|---------|
| SEARCHED | INDEXED |
| SERIALIZED | FILED |
| APR 1982 | |
| FBI - BALTIMORE | |
| A | |

Blank

CONTENTS

| | Page |
|---|------|
| 1. INTRODUCTION | 7 |
| 2. BLANDAMER'S METHOD FOR COMPUTING ACTIVATION PARAMETERS. | 8 |
| 3. RESULTS AND DISCUSSION | 9 |
| 4. CONCLUSIONS | 9 |
| LITERATURE CITED | 11 |
| APPENDIXES | |
| Appendix A - Collected Tables | 15 |
| Appendix B - Temperature Fit of Formation Constant at Various Temperatures | 23 |
| DISTRIBUTION LIST | 31 |


Blank

ON THE TEMPERATURE DEPENDANCE OF THE FORMATION CONSTANT OF THIOCYANATOPENTAQUOCHROMIUM (III) IN ACIDIC SOLUTION

1. INTRODUCTION

The ionization constants for the reversible dissociation of carboxylic acids pass through a minimum as temperature varies.¹ If the ionization is regarded as a one-step or unitary process, then the heat capacity at constant pressure, ΔC_p° , must be negative. Numerous investigators have tried to find expressions to fit the experimentally observed temperature dependence in order to compute ΔC_p° which is regarded as a measure of solute-solvent interactions.² Timimi reviewed these empirical expressions.³

More recently, Blandamer⁴ and coworkers in the United Kingdom and Canada have reexamined the interpretation of negative heat capacities for the ionization of carboxylic acids, using Alberty and Robinson's idea⁵ that such nonzero heat capacities are artifacts caused by the presence of intermediates. Starting with a suggestion by Eigen⁶ that diffusion-controlled intermediates are formed in the ionization of carboxylic acids, Blandamer and coworkers postulated^{7,8} that the dissociation of carboxylic acids could be written as



where the intermediates are hydrogen-bonded, diffusion-controlled, "encounter" complexes with K_1 and K_3 values near unity. Since $K_2 \ll 1$ for carboxylic acids, the observed dissociation constant, K_0 , is


$$K_0 = K_1 K_2 K_3 / (1 + K_1) \quad (2)$$

Blandamer showed that the observed temperature dependence for K_0 arose from equation (2) using temperature-independent enthalpies for the individual equilibria K_1 , K_2 , and K_3 .

Over the past three years, Blandamer⁹⁻¹⁷ has also shown that the solvolysis of many organic compounds with nonzero heat capacities of activation which had been treated as unitary reactions¹⁸ may also be multistep reactions with rapid preequilibria. Blandamer and colleagues have labeled such temperature-dependent activation energies, arising from the presence of preequilibria, as "spurious" or "anomalous." Blandamer noted that spurious heat capacities are themselves temperature-dependent,^{5,19} and an expression that he derived to compute enthalpies and heat capacities of activation also gave temperature-dependent ΔC_p° , if preequilibria were present.¹³ Alberty and Robinson⁵ chided earlier investigators for fitting rate or equilibrium constants to expressions with temperature-independent heat capacities.

In contrast to the wealth of information on organic compounds with temperature-dependent activation energies, there are far fewer examples in the inorganic

chemistry literature, particularly for formation constants of strong complexes which one can liken to the formation of carboxylic acids (reverse of equation 2).²⁰ A notable exception is the formation of thiocyanatopentaaquochromium (III), given below:



Postmus and King^{21, 22} found the equilibrium constant for equation 3 passed through a minimum in accordance with the behavior of the observed association constant for carboxylic acids. Postmus and King computed a value of ΔC_p° for equation 3, but they used an expression with a temperature-independent heat capacity.

In view of Blandamer's recent interpretation of carboxylic acid ionization, one wonders whether the ΔC_p° that Postmus and King measured is also "spurious." To this end, Blandamer's expression has been applied to Postmus and King's equilibrium constants to test the temperature-dependence of ΔC_p° for equation 3.

2. BLANDAMER'S METHOD FOR COMPUTING ACTIVATION PARAMETERS

Blandamer's expression for calculation of thermodynamic parameters from the temperature dependence of equilibrium data is

$$K = K_0 e \left\{ \frac{\Delta H^\circ}{R} \left(\frac{1}{T_0} - \frac{1}{T} \right) + \frac{\Delta C_p^\circ}{R} \left[\ln \left(\frac{T}{T_0} \right) + \frac{T}{T_0} + 1 \right] \right\} \quad (4)$$

where K = equilibrium constant at temperature, T
 K_0 = equilibrium constant at temperature, T_0
 ΔH° = enthalpy at T_0 , and
 ΔC_p° = heat capacity at T_0 .

The equation is obtained by the integration of the van't Hoff isochore between the temperature T_0 and T assuming ΔC_p° is independent of temperature. If ΔC_p° is dependent on temperature, equation 4 gives a poor fit and the ΔC_p° values as computed do predict the correct trend of the actual ΔC_p° dependence with temperature, though thermodynamically incorrect. For rate data, K and K_0 in equation 4 become k and $(k_0 \cdot T)/T_0$, respectively, and ΔH° and ΔC_p° become activation parameters. For a series of measurements at n pairs of K and T , any pair of K and T is set as K_0 and T_0 , and all other values of K, T are fit to equation 4 to give ΔH° and ΔC_p° at Temperature T_0 . The calculation is repeated with a different K, T set as K_0, T_0 , until all values of T have been used as T_0 . This results in n values of ΔH° and ΔC_p° from which one determines whether ΔC_p° is zero, constant, or temperature dependent.

A nonlinear least-squares program²³ is used to find the best-fit values of ΔH° and ΔC_p° from a set of K, T data. The temperature, T , is the independent variable, K is the dependent variable, ΔH° and ΔC_p° are parameters to be fit, and K_0 and T_0 are fixed parameters. The program finds best-fit values of ΔH° and ΔC_p° and then computes the equilibrium constants with these best-fit values of ΔH° and ΔC_p° to compare the fit between sets of data.

3. RESULTS AND DISCUSSION

Table A-1* lists the equilibrium constants Postmus and King measured for the formation constant of $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$, which were corrected to zero ionic strength with a modified Debye-Huckel equation. Three formation constants at 14.0°C, 25.1°C and 30.0°C are the ratio of the forward and reverse rate constants. At 30°C, the formation constant was measured both spectrophotometrically and as a ratio of rate constants.

In table A-2, which illustrates typical results of the temperature fit with Blandamer's expression using 94.6°C as an example, one can see that the formation constants calculated with the best-fit values of ΔH° and ΔC_p° agree well with the experimental values. The results of the calculations at each of the other temperatures are listed in appendix B. Table A-3 summarizes the values of ΔH° and ΔC_p° from all the calculations, which seem to show that ΔC_p° is constant. To test this premise further, the enthalpies in table A-3 were fit linearly with temperature with the least-squares program. The results of this calculation are shown in table A-4. Again, agreement between experimental and computed enthalpies and small standard deviations of the mean suggest that the fit is adequate and that ΔC_p° can be considered constant at 66 cal/mole-K.

These results suggest, then, that the ΔC_p° for the formation constant of $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$ is not spurious, and that temperature-dependent enthalpies should be much more common for inorganic reactions, one need only extend the temperature range to find the temperature dependence. As table A-5 shows, many inorganic equilibria²⁴⁻²⁷ have been measured over narrow temperature ranges, in contrast to Postmus and King's experiments. In all instances in table A-6 where four temperature measurements are available, application of Blandamer's method (equation 4) revealed temperature-dependent enthalpies (tables A-6 and A-7). Postmus and King²¹ also noted that the acid dissociation constant of $\text{Cr}(\text{H}_2\text{O})_6^{3+}$ had a temperature-dependent ΔH° , though much smaller than that for the formation constant of $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$.

These results also caution against extrapolating rate or equilibrium data beyond the measured temperature range. Assume, for example, that one used only the three formation constants from the ratio of rate constants to compute ΔH° by the following:

$$K = e^{(-\Delta H^\circ/RT + \Delta S^\circ/R)} \quad (5)$$

Table 8 shows the fit obtained with the least-squares program. One would conclude that the fit is adequate over this temperature range, but it would be a mistake to extrapolate to 94.6°C with this value of ΔH° .

4. CONCLUSIONS

a. The heat capacity for the formation constant of $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$ is constant with temperature. This suggests that the heat capacity is not spurious, in the sense of arising from establishment of rapid preequilibria.

*See appendix A for tables.

b. The formation constants of many inorganic equilibria should also have temperature-dependent enthalpies. Failure to see such temperature dependences may rest on the narrow temperature range over which most measurements have been made in comparison to Postmus and King's measurement with $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$.

LITERATURE CITED

1. Lewis, G. N., and Randall, M. *Thermodynamics*. 2nd Edition. Revised by Pitzer, K. S. and Brewer, L. Chapter 22. p 305. McGraw-Hill Book Co., New York, New York, 1961.
2. Kohnstam, G. Heat Capacities of Activation and Their Uses in Mechanistic Studies. *Advances in Physical Organic Chemistry*. 5, 122 to 172 (1967).
3. Timimi, B. A. The Evaluation of ΔC_p° for Acid-Base Equilibria from pK Measurement. *Electrochimica Acta*. 19, 149 to 158 (1974).
4. Blandamer, M. J., Scott, J. W. M., and Robertson, R. E. Analysis of the Dependence of Acid Dissociation Constants on Temperature. *J. Chem. Soc. Perkin II*, 447 to 456 (1981).
5. Albery, W. J. and Robinson, B. H. Measurement of Enthalpies and Heat Capacities of Activation by Differential Methods. *J. Chem. Soc., Trans. Faraday Soc.* 65, 980 to 991 (1969).
6. Eigen, M. Proton Transfer, Acid-Base Catalysis, and Enzymatic Hydrolysis. Part I: Elementary Processes. *Angewandte Chemie, Int'l Ed.* 3, 1 to 72 (1964).
7. Blandamer, M. J., Burgess, J., Duce, P. P., Robertson, R. E., and Scott, J. M. W. Analysis of the Dependence on Temperature of the Acid Dissociation Constants for Mono-Carboxylic Acids in Water in Terms of a Two-Stage Mechanism. *J. Chem. Soc., Trans. Faraday Soc. I.* 77, 2281 to 2286 (1981).
8. Blandamer, M. J., Robertson, R. E., and Scott, J. M. W. Thermodynamics of Acid-Base Reactions in Water: A Comment on the Eigen Mechanism. *J. Am. Chem. Soc.* 104, 1136 to 1138 (1982).
9. Blandamer, M. J., Robertson, R. E., and Scott, J. M. W. An Examination of Parameters Describing the Temperature Dependence of Rate Constants on temperature for Solvolysis of Various Organic Esters in Water and Aqueous Mixtures. *Can. J. Chem.* 58, 772 to 776 (1980).
10. Blandamer, M. J., Robertson, R. E., Scott, J. M. W., and Vrieling, A. Evidence for the Incursion of Intermediates in the Hydrolysis of Tertiary, Secondary, and Primary Substrates. *J. Am. Chem. Soc.* 102, 2585 to 2592 (1980).
11. Blandamer, M. J., Burgess, J., Duce, P. P., Robertson, R. E., and Scott, J. M. W. The Dependence on Temperature of Rate Constants for the Solvolysis of Alkyl Halides and Related Compounds in Water. *J. Chem. Soc. Perkin II*, 1157 to 1160 (1981).
12. Blandamer, M. J., Burgess, J., Duce, P. P., Robertson, R. E., and Scott, J. M. W. A Re-Examination of the Effects of Added Solvent on the Activation Parameters for Solvolysis of t-Butyl Chloride in Water. *J. Chem. Soc., Trans. Faraday Soc. I.* 77, 1999 to 2008 (1981).

13. Blandamer, M. J., Robertson, R. E., Golding, P. D., MacNeil, J. M., and Scott, J. M. W. On the Heat Capacities of Activation for Displacements at Primary and Secondary Carbon Centers in Water. *J. Am. Chem. Soc.* 103, 2415 to 2416 (1981).
14. Blandamer, M. J., Burgess, J., Duce, P. P., Robertson, R. E., and Scott, J. M. W. The Dependence of Acid Dissociation Constants in Water on Temperature as Expressed by the Guinney Equation. *Can. J. Chem.* 59, 2845 to 2847 (1981).
15. Blandamer, M. J., Robertson, R. E., Scott, J. M. W., and Golding, P. E. Some Observations Concerning the Temperature Dependence of the Reaction of Ammonia with Water. *J. Am. Chem. Soc.* 103, 5923 to 5925 (1981).
16. Blandamer, M. J., Burgess, J., Duce, P. P., Robertson, R. E., and Scott, J. M. W. Effects of Added Acetonitrile on the Heat Capacities of Activation for the Solvolysis of Sample Organic Esters in Water. *J. Chem. Soc., Faraday Trans. I.* 78, 881 to 886 (1982).
17. Blandamer, M. J., Burgess, J., Clare, N. P., Duce, P. P., Gray, R. P., Robertson, R. E., and Scott, J. M. W. Analysis of the Dependence on Temperature of Kinetic Solvent Isotope Effects. *J. Chem. Soc., Faraday Trans. I.* 78, 1103 to 1115 (1982).
18. Blandamer, M. J. Kinetics of Organic Reactions in Water and Aqueous Mixtures. *Advances in Physical Organic Chemistry.* 14, 203 to 352 (1976).
19. Koren, R. and Perlmutter-Hayman. Rapid Pre-equilibria as a Cause for Change of Activation Energy with Temperature. *J. Phys. Chem.* 75, 2372 to 2376 (1971).
20. Rossitti, F. J. C. "The Thermodynamics of Metal Ion Complex Formation in Solution." *Modern Coordination Chemistry=Principles and Methods*, Lewis, J. and Wilkins, R. G., eds. Chapter 1. Interscience Publishers Inc., New York. New York. 1960.
21. Postmus, C. and King, E. L. The Equilibria in Acidic Solutions of Chromium (III) Ion and Thiocyanate Ion. *J. Phys. Chem.* 59, 1208 to 1216 (1955).
22. Postmus, C., and King, E. L. The Rate Law for the Forward and Reverse of the Reaction $\text{Cr}(\text{H}_2\text{O})_6^{3+} + \text{NCS}^- \rightleftharpoons \text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+} + \text{H}_2\text{O}$. *J. Phys. Chem.* 59, 1216 to 1221 (1955).
23. Moore, R. H., and Ziegler, R. K. Los Alamos Scientific Laboratory Report LA-2367. The Solution of the General Least-Squares Problem with Special Reference to High-Speed Computers. March 1960. UNCLASSIFIED Report.
24. Betts, R. H. and Dainton, F. S. Electron Transfer and Other processes Involved in the Spontaneous Bleaching of Acidified Aqueous Solutions of Ferric Thiocyanate. *J. Am. Chem. Soc.* 75, 5721-5727 (1953).
25. Milburn, R. N. A Spectrophotometric Study of the Hydrolysis of Iron (III). III. Heats and Entropies of Hydrolysis. *J. Am. Chem. Soc.* 79, 537 to 540 (1957).

26. Sutcliffe, L. H., and Weber, J. R. The Reaction Between Cerium (III) and Cobalt (III). J. Chem. Soc., Trans. Faraday Soc. 52, 1225 to 1234 (1956).

27. Hardwicke, T. J. and Robertson, E. Ionic Species in Ceric Perchlorate Solutions. Can. J. Chem. 29, 818 to 827 (1951).

Blank

APPENDIX A
COLLECTED TABLES

Table A-1. Formation Constants for $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$ ^a

| K $\text{M}^{-1} \times 10^{-3}$ | T $^{\circ}\text{C}$ |
|---------------------------------------|---------------------------|
| 1.24 | 94.6 |
| 1.15 | 84.8 |
| 1.06 | 73.7 |
| 1.04 | 63.6 |
| 1.03 | 46.2 |
| 1.10 | 30.0 |
| 1.12 ^b | 30.0 |
| 1.23 ^b | 25.1 |
| 1.38 ^b | 14.0 |

a Taken from reference 21. Formation constants corrected to zero ionic strength.

b Ratio of forward and reverse rate constants (reference 22).

Table A-2. Temperature Fit of Formation Constant at 94.6° C

| T | K, experimental | K, calculated* |
|-------------------|------------------------------------|------------------------------------|
| ^o C | M ⁻¹ x 10 ⁻³ | M ⁻¹ x 10 ⁻³ |
| 84.8 | 1.15 | 1.14 |
| 73.7 | 1.06 | 1.06 |
| 63.6 | 1.04 | 1.03 |
| 46.2 | 1.03 | 1.03 |
| 30.0 | 1.10 | 1.13 |
| 30.0 | 1.12 | 1.13 |
| 25.1 | 1.23 | 1.19 |
| 14.0 | 1.38 | 1.38 |

* $\Delta H^{\circ} = 2,535 \pm 150$ cal/mole; $\Delta C_p^{\circ} = 64.7 \pm 4$; error expressed as standard deviation of the mean.

Table A-3. Summary of Enthalpies and Heat Capacities
for Formation Constant of $\text{Cr}(\text{H}_2\text{O})_5\text{NCS}^{2+}$

| T | ΔH° | ΔC_p° |
|------------------|-------------------|--------------------|
| $^\circ\text{C}$ | cal/mole | cal/mole-K |
| 94.6 | $2,535 \pm 150^*$ | $64.7 \pm 4^*$ |
| 84.8 | $1,991 \pm 169$ | 66.6 ± 5 |
| 73.7 | $1,201 \pm 182$ | 66.2 ± 7 |
| 63.6 | 548 ± 132 | 62.6 ± 6 |
| 46.2 | -580 ± 57 | 65.7 ± 4 |
| 30.0 | $-1,765 \pm 183$ | 76.6 ± 8 |
| 30.0 | $-1,693 \pm 126$ | 69.9 ± 8 |
| 25.1 | $-1,998 \pm 275$ | 61.1 ± 12 |
| 14.0 | $-2,668 \pm 135$ | 65.1 ± 5 |

*Error expressed as standard deviation of the mean.

Table A-4. Linear Fit of Enthalpy versus Temperature

| T °C | ΔH° cal/mole | ΔH° calculated* cal/mole |
|---------|------------------------------|--|
| 94.6 | 2,535 | 2,579 |
| 84.8 | 1,991 | 1,934 |
| 73.7 | 1,201 | 1,203 |
| 63.6 | 548 | 9,538 |
| 46.2 | -580 | -608 |
| 30.0 | -1,765 | -1,675 |
| 30.0 | -1,693 | -1,675 |
| 25.1 | -1,998 | -1,998 |
| 14.0 | -2,668 | -2,728 |

* $\Delta H^\circ = -3,650 \pm 36$ cal/mole; $\Delta C_p^\circ = 65.9 \pm 0.6$ cal/mole-K

Table A-5. Temperature Range for Measuring Some Inorganic Equilibria

| Reaction | No. of Measurements | Range | Reference |
|---|---------------------|--------------|-----------|
| | | °C | |
| $\text{Fe} + \text{NCS} \rightleftharpoons \text{FeNCS}^{2+}$ | 3 | 11.4 to 28.5 | 24 |
| $\text{Fe}^{3+} \rightleftharpoons \text{FeOH}^{2+} + \text{H}^+$ | 3 | 18 to 32 | 25 |
| $2\text{FeOH}^{2+} \rightleftharpoons (\text{FeOH})_2^{4+}$ | 3 | 18 to 32 | 25 |
| $\text{Co}^{3+} \rightleftharpoons \text{CoOH}^{2+} + \text{H}^+$ | 4 | 12.5 to 28.2 | 26 |
| $\text{Ce}^{4+} \rightleftharpoons \text{CeOH}^{3+} + \text{H}^+$ | 4 | 5 to 35 | 27 |
| $2\text{CeOH}^{3+} \rightleftharpoons [\text{Ce-O-Ce}]^{6+} + \text{H}_2\text{O}$ | 4 | 5 to 35 | 27 |

Table A-6. Enthalpy and Heat Capacity
for Cobaltic Ion Dissociation
Using Equation (4)

| T | ΔH° | ΔC_p° |
|------------------|------------------|--------------------|
| $^\circ\text{C}$ | kcal/mole | kcal/mole-K |
| 12.5 | 10.3 | -2.0 |
| 18.4 | 10.1 | -2.0 |
| 23.6 | 2.0 | -2.0 |
| 28.2 | 1.5 | -0.7 |



Table A-7. Enthalpy and Heat Capacities by Equation (4)
for Ceric Ion Dissociation^a and Dimerization^b

| Temperature | $\Delta H^{\circ a}$ | $\Delta C_p^{\circ a}$ | $\Delta H^{\circ b}$ | $\Delta C_p^{\circ b}$ |
|-------------|----------------------|------------------------|----------------------|------------------------|
| °C | kcal/mole | kcal/mole-K | kcal/mole-K | kcal/mole-K |
| 5 | 11.8 | 0.30 | 42.4 | -2.9 |
| 15 | 13.6 | 0.39 | -10.8 | -1.1 |
| 25 | 17.4 | 0.39 | -19.5 | -0.7 |
| 35 | 21.3 | 0.38 | -31.3 | -0.8 |





Table A-8. Temperature Fit of Formation Constant Using Kinetic Data

| Temperature | K, experimental | K, fit* |
|-------------|-------------------------|-------------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ |
| 30.0 | 1.10 | 1.12 |
| 25.1 | 1.23 | 1.20 |
| 14.0 | 1.38 | 1.39 |

* $\Delta H = -2.3 \pm 0.5$ kcal/mole; $\Delta S = 6.5 \pm 1.7$ cal/mole-K.

APPENDIX B

TEMPERATURE FIT OF FORMATION CONSTANT AT VARIOUS TEMPERATURES

Table B-1. Temperature Fit of Formation Constant at 84.8°C

| Temperature | K, experimental | K, calculated * |
|-------------|-------------------------|-------------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ |
| 94.6 | 1.24 | 1.25 |
| 73.7 | 1.06 | 1.07 |
| 63.6 | 1.04 | 1.03 |
| 46.2 | 1.03 | 1.03 |
| 30.0 | 1.10 | 1.13 |
| 30.0 | 1.12 | 1.13 |
| 25.1 | 1.23 | 1.19 |
| 14.0 | 1.38 | 1.36 |

* $\Delta H^\circ = 1,991 \pm 169$ cal/mole; $\Delta C_p^\circ = 66.6 \pm 5$ cal/mole·K.

Table B-2. Temperature Fit of Formation Constant at 73.7°C

| Temperature | K, experimental | K, calculated* |
|-------------|-------------------------|-------------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ |
| 94.6 | 1.24 | 1.24 |
| 84.8 | 1.15 | 1.14 |
| 63.6 | 1.04 | 1.02 |
| 46.2 | 1.03 | 1.02 |
| 30.0 | 1.10 | 1.13 |
| 30.0 | 1.12 | 1.13 |
| 25.1 | 1.23 | 1.19 |
| 14.0 | 1.38 | 1.39 |

* $\Delta H^\circ = 1201 \pm 182$ cal/mole; $\Delta C_p^\circ = 66.2 \pm 7$ cal/mole-K.

Table B-3. Temperature Fit of Formation Constant at 63.6°C

| Temperature °C | K, experimental $M^{-1} \times 10^{-3}$ | K, calculated* $M^{-1} \times 10^{-3}$ |
|-------------------|--|---|
| 94.6 | 1.24 | 1.25 |
| 84.8 | 1.15 | 1.16 |
| 73.7 | 1.06 | 1.08 |
| 46.2 | 1.03 | 1.04 |
| 30.0 | 1.10 | 1.14 |
| 30.0 | 1.12 | 1.14 |
| 25.1 | 1.23 | 1.19 |
| 14.0 | 1.38 | 1.38 |

* $\Delta H^\circ = 548 \pm 132$ cal/mole; $\Delta C_p^\circ = 62.6 \pm 6$ cal/mole-K.

Table B-4. Temperature Fit of Formation Constant at 46.2°C

| Temperature °C | K, experimental $M^{-1} \times 10^{-3}$ | K, calculated * $M^{-1} \times 10^{-3}$ |
|-------------------|--|--|
| 94.6 | 1.24 | 1.25 |
| 84.8 | 1.15 | 1.15 |
| 73.7 | 1.06 | 1.07 |
| 63.6 | 1.04 | 1.03 |
| 30.0 | 1.10 | 1.13 |
| 30.0 | 1.12 | 1.13 |
| 25.1 | 1.23 | 1.19 |
| 14.0 | 1.38 | 1.38 |

* $\Delta H^\circ = -580 \pm 57$ cal/mole; $\Delta C_p^\circ = 65.7 \pm$ cal/mole-K.

Table B-5. Temperature Fit of Formation Constant at 30.0°C

| Temperature | K, experimental | K, calculated | |
|-------------|-------------------------|-------------------------|-------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ | |
| 94.6 | 1.24 | 1.29 ^a | 1.27 ^b |
| 84.8 | 1.15 | 1.16 | 1.15 |
| 73.7 | 1.06 | 1.06 | 1.07 |
| 63.6 | 1.04 | 1.01 | 1.02 |
| 46.2 | 1.03 | 1.00 | 1.02 |
| 25.1 | 1.23 | 1.16 | 1.18 |
| 14.0 | 1.38 | 1.37 | 1.38 |

a $\Delta H^{\circ} = -1,765 \pm 183 \text{ cal/mole}$; $\Delta C_p^{\circ} = 76.6 \pm 8 \text{ cal/mole-K}$
 $(K_{30.0^{\circ}\text{C}} = 1.10 \times 10^3 M^{-1})$

* b $\Delta H^{\circ} = 1,693 \pm 126 \text{ cal/mole}$; $\Delta C_p^{\circ} = 69.9 \pm 6 \text{ cal/mole-K}$
 $(K_{30.0^{\circ}\text{C}} = 1.12 \times 10^3 M^{-1})$

Table B-6. Temperature Fit of Formation Constant at 25.1°C

| Temperature | K, experimental | K, calculated* |
|-------------|-------------------------|-------------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ |
| 94.6 | 1.24 | 1.22 |
| 84.8 | 1.15 | 1.14 |
| 73.7 | 1.06 | 1.07 |
| 63.6 | 1.04 | 1.04 |
| 46.2 | 1.03 | 1.06 |
| 30.0 | 1.10 | 1.17 |
| 30.0 | 1.12 | 1.17 |
| 14.0 | 1.38 | 1.43 |

* $\Delta H^\circ = -1998 \pm 275$ cal/mole; $\Delta C_p^\circ = 61.1 \pm 12$ cal/mole-K.

Table B-7. Temperature Fit of Formation Constant at 14.0°C

| Temperature | K, experimental | K, calculated* |
|-------------|-------------------------|-------------------------|
| °C | $M^{-1} \times 10^{-3}$ | $M^{-1} \times 10^{-3}$ |
| 94.6 | 1.24 | 1.25 |
| 84.8 | 1.15 | 1.15 |
| 73.7 | 1.06 | 1.07 |
| 63.6 | 1.04 | 1.03 |
| 46.2 | 1.03 | 1.03 |
| 30.0 | 1.10 | 1.13 |
| 30.0 | 1.12 | 1.13 |
| 25.1 | 1.23 | 1.19 |

* $\Delta H^\circ = -2668 \pm 135$ cal/mole; $\Delta C_p^\circ = 65.1 \pm 5$ cal/mole-K.

Blank

DISTRIBUTION LIST 2

| Names | Copies | Names | Copies |
|--------------------------------------|--------|---|--------|
| CHEMICAL SYSTEMS LABORATORY | | Federal Emergency Management Agency | |
| ATTN: DRDAR-CLB | 1 | Office of Research/NPP | |
| ATTN: DRDAR-CLB-C | 1 | ATTN: David W. Bensen | 1 |
| ATTN: DRDAR-CLB-PO | 1 | Washington, DC 20472 | |
| ATTN: DRDAR-CLB-R | 1 | | |
| ATTN: DRDAR-CLB-R(M) | 1 | HQ DA | |
| ATTN: DRDAR-CLB-R(S) | 1 | Office of the Deputy Chief of Staff for | |
| ATTN: DRDAR-CLB-T | 1 | Research, Development & Acquisition | |
| ATTN: DRDAR-CLC-B | 1 | ATTN: DAMA-CSS-C | 1 |
| ATTN: DRDAR-CLC-C | 1 | Washington, DC 20310 | |
| ATTN: DRDAR-CLC-E | 1 | | |
| ATTN: DRDAR-CLF | 1 | HQ Sixth US Army | |
| ATTN: DRDAR-CLJ-R | 1 | ATTN: AFKC-OP-NBC | 1 |
| ATTN: DRDAR-CLJ-L | 2 | Presidio of San Francisco, CA 94129 | |
| ATTN: DRDAR-CLJ-M | 1 | | |
| ATTN: DRDAR-CLN | 1 | Commander | |
| ATTN: DRDAR-CLT | 1 | DARCOM, STITEUR | |
| ATTN: DRDAR-CLW-C | 1 | ATTN: DRXST-STI | 1 |
| ATTN: DRDAR-CLW-P | 1 | Box 48, APO New York 09710 | |
| ATTN: DRDAR-CLY-A | 1 | | |
| ATTN: DRDAR-CLY-R | 4 | Commander | |
| | | USASTCFEO | |
| COPIES FOR AUTHOR(S) | | ATTN: MAJ Mikeworth | 1 |
| Research Division | 6 | APC San Francisco 96328 | |
| RECORD COPY: DRDAR-CLB-A | 1 | | |
| | | Commander | |
| DEPARTMENT OF DEFENSE | | US Army Nuclear & Chemical Agency | |
| | | ATTN: MONA-WE | 1 |
| Defense Technical Information Center | 12 | 7500 Backlick Rd, Bldg 2073 | |
| ATTN: DTIC-DDA-2 | | Springfield, VA 22150 | |
| Cameron Station, Building 5 | | | |
| Alexandria, VA 22314 | | Army Research Office | |
| | | ATTN: DRXRO-CB (Dr. R. Ghirardelli) | 1 |
| Director | | P.O. Box 12211 | |
| Defense Intelligence Agency | | Research Triangle Park, NC 27709 | |
| ATTN: DB-4G1 | 1 | | |
| Washington, DC 20301 | | OFFICE OF THE SURGEON GENERAL | |
| | | Commander | |
| Commander | | US Army Medical Bioengineering Research | |
| USASED, USAINSCOM | | and Development Laboratory | |
| ATTN: IAFM-SED-III | 1 | ATTN: SGRD-UBD-AL, Bldg 568 | 1 |
| Fort Meade, MD 20755 | | Fort Detrick, Frederick, MD 21701 | |
| | | | |
| DEPARTMENT OF THE ARMY | | Commander | |
| HQDA | | USA Medical Research Institute of | |
| ATTN: DAMO-NCC | 1 | Chemical Defense | |
| ATTN: DAMO-NC/COL Robinson (P) | 1 | ATTN: SGRD-UV-L | 1 |
| WASH DC 20310 | | Aberdeen Proving Ground, MD 21010 | |

US ARMY MATERIEL DEVELOPMENT AND
READINESS COMMAND

Commander

US Army Materiel Development and
Readiness Command

ATTN: DRCLDC 1
ATTN: DRCSF-P 1
5001 Eisenhower Ave
Alexandria, VA 22333

Project Manager Smoke/Obscurants

ATTN: DRCPM-SMK-S 3
Aberdeen Proving Ground, MD 21005

Commander

US Army Foreign Science & Technology Center
ATTN: DRXST-MT3 1
220 Seventh St., NE
Charlottesville, VA 22901

Director

US Army Materiel Systems Analysis Activity
ATTN: DRXSY-MP 1
ATTN: DRXSY-CA (Mr. Metz) 1
Aberdeen Proving Ground, MD 21005

Commander

US Army Missile Command
Redstone Scientific Information Center
ATTN: DRSMI-RPR (Documents) 1
Redstone Arsenal, AL 35809

Director

DARCOM Field Safety Activity
ATTN: DRXOS-C 1
Charlestown, IN 47111

Commander

US Army Natick Research and Development
Laboratories

ATTN: DRDNA-O 1
ATTN: DRDNA-IC 1
ATTN: DRDNA-IM 1
ATTN: DRDNA-ITF (Dr. Roy W. Roth) 2
Natick, MA 01760

US ARMY ARMAMENT RESEARCH AND
DEVELOPMENT COMMAND

Commander

US Army Armament Research and
Development Command

ATTN: DRDAR-LCA-L 1
ATTN: DRDAR-LCE-C 1
ATTN: DRDAR-LCU-CE 1
ATTN: DRDAR-NC (COL Lymn) 3
ATTN: DRDAR-SCA-T 1
ATTN: DRDAR-SCM 1
ATTN: DRDAR-SCP 1
ATTN: DRDAR-SCS 1
ATTN: DRDAR-TDC (Dr. D. Gyorog) 1
ATTN: DRDAR-TSS 2
ATTN: DRCPM-CAWS-AM 1
Dover, NJ 07801

US Army Armament Research and
Development Command

Resident Operations Office

ATTN: DRDAR-TSE-OA (Robert Thresher) 1
National Space Technology Laboratories
NSTL Station, Mississippi 39529

Commander

ARRADCOM
ATTN: DRDAR-QAC-E 1
Aberdeen Proving Ground, MD 21010

Commander

USA Technical Detachment 1
US Naval EOD Technology Center
Indian Head, MD 20640

US ARMY ARMAMENT MATERIEL READINESS
COMMAND

Commander

US Army Armament Materiel Readiness Command

ATTN: DRSAR-ASN 1
ATTN: DRSAR-IRW 1
Rock Island, IL 61299

Commander

US Army Dugway Proving Ground
ATTN: Technical Library (Docu Sect) 1
Dugway, UT 84022

US ARMY TRAINING & DOCTRINE COMMAND

Commandant

US Army Infantry School
ATTN: CTDD, CSD, NBC Branch
Fort Benning, GA 31905

1

Commandant

US Army Missile & Munitions Center
and School
ATTN: ATSK-CM
ATTN: ATSK-TME
Redstone Arsenal, AL 35809

1

1

Commander

US Army Logistics Center
ATTN: ATCL-MG
Fort Lee, VA 23801

1

Commandant

US Army Chemical School
ATTN: ATZN-CM-C
ATTN: ATZN-CM-AD
ATTN: ATZN-CM-TPC
Fort McClellan, AL 36205

1

2

2

Commander

USAAVNC
ATTN: ATZQ-D-MS
Fort Rucker, AL 36362

1

Commander

US Army Infantry Center
ATTN: ATSH-CD-MS-C
Fort Benning, GA 31905

1

Commander

USA Training and Doctrine Command
ATTN: ATCD-N
Fort Monroe, VA 23651

1

Commander

US Army Armor Center
ATTN: ATZK-CD-MS
ATTN: ATZK-PPT-PO-C
Fort Knox, KY 40121

1

1

Commander

USA Combined Arms Center and
Fort Leavenworth
ATTN: ATZL-CAM-IM
Fort Leavenworth, KS 66027

1

US ARMY TEST & EVALUATION COMMAND

Commander

US Army Test & Evaluation Command
ATTN: DRSTE-CT-T
Aberdeen Proving Ground, MD 21005

1

DEPARTMENT OF THE NAVY

Chief of Naval Research

ATTN: Code 441
800 N. Quincy Street
Arlington, VA 22217

1

Project Manager

Theatre Nuclear Warfare Project Office
ATTN: TN-09C
Navy Department
Washington, DC 20360

1

Commander

Naval Explosive Ordnance Disposal
Technology Center
ATTN: AC-3
Indian Head, MD 20640

1

Commander

Naval Surface Weapons Center
Code G51
Dahlgren, VA 22448

1

Chief, Bureau of Medicine & Surgery Department of the Navy

ATTN: MED 3C33
Washington, DC 20372

1

Commander

Naval Air Development Center
ATTN: Code 2012 (Dr. Robert Helmbold)
Warminster, PA 18974

1

US MARINE CORPS

Commandant

HQ, US Marine Corps
ATTN: Code LMW-50
Washington, DC 20380

1

Commanding General

Marine Corps Development and
Education Command
ATTN: Fire Power Division, D091
Quantico, VA 22134

1

DEPARTMENT OF THE AIR FORCE

ASD/AESD
Wright-Patterson AFB, OH 45433

HQ AFSC/SDZ
ATTN: CPT D. Riediger
Andrews AFB, MD 20334

HQ, AFSC/SDNE
Andrews AFB, MD 20334

HQ, AFSC/SGB
Andrews AFB, DC 20334

HQ, NORAD
ATTN: J-3TU
Peterson AFB, CO 80914

AFAMRL/HE
ATTN: Dr. Clyde Reploggle
Wright-Patterson AFB, OH 45433

HQ AFTEC/TEL
Kirtland AFB, NM 87117

USAF TAWC/THL
Eglin AFB, FL 32542

AFATL/DLV
Eglin AFB, FL 32542

USAF SC
ATTN: AD/YQ
ATTN: AD/YQO (MAJ Owens)
Eglin AFB, FL 32542

USAFSAM/VN
Deputy for Chemical Defense
ATTN: Dr. F. Wesley Baumgardner
Brooks AFB, TX 78235

AFAMRL/TS
ATTN: COL Johnson
Wright-Patterson AFB, OH 45433

AMD/RDTK
ATTN: LTC T. Kingery
Brooks AFB, TX 78235

AMD/RDSM
Brooks AFB, TX 78235

AMD/RDSX
Brooks AFB, TX 78235

AD/XRO
Eglin AFB, FL 32542

OUTSIDE AGENCIES

Battelle, Columbus Laboratories
ATTN: TACTEC
505 King Avenue
Columbus, OH 43201

Toxicology Information Center, JH 652
National Research Council
2101 Constitution Ave., NW
Washington, DC 20418

US Public Health Service
Center for Disease Control
ATTN: Lewis Webb, Jr.
Building 4, Room 232
Atlanta, GA 30333

Director
Central Intelligence Agency
ATTN: AMR/ORD/DD/S&T
Washington, DC 20505

ADDITIONAL ADDRESSEE

Commandant
Academy of Health Sciences, US Army
ATTN: HSHA-CDH
ATTN: HSHA-IPM
Fort Sam Houston, TX 78234

END

FILMED

6-83

DTIC